

Staged Surgical Intervention in the Treatment of Septic Ankle Arthritis with Autologous Circular Pillar Fibula Augmentation: A Case Report

Sham J. Persaud DPM; Alan R. Catanzariti, DPM, FACFAS

The Foot & Ankle Institute, West Penn Hospital, Allegheny Health Network, Pittsburgh, PA

Purpose

Surgical management of chronic septic arthritis of the ankle joint is a challenging problem. Failure to initiate appropriate antibiotic therapy and perform I&D within the first 24 to 48 hours of onset can result in subchondral bone loss and permanent joint dysfunction. Literature states joint function after *S. aureus* septic arthritis is generally 46-50% of its baseline. Cierny et al. related a 25% amputation rate for patients with fusions of septic ankle joints. This case study presents a staged procedure for limb salvage of a patient with chronic septic arthritis of the ankle joint.

Case Study

A 54-year-old female with chronic right septic ankle arthritis for 6 months presented for evaluation. The patient had undergone arthrocentesis with corticosteroid, as well as incision and drainage with washout and long term IV antibiotic therapy. She was offered a below knee amputation but was reluctant to proceed and sought a second opinion. Her pre-operative imaging is seen in Figure 1 A-B. The patient chose to proceed with a staged surgical approach for limb salvage.

The patient underwent needle biopsy of the tibia and talus with arthroscopic debridement. Arthroscopy was performed in standard fashion using a 2.7mm 30 degree arthroscope, utilizing a burr and shaver for ankle joint debridement. Arthroscopic evaluation of the ankle joint revealed destruction of both tibial and talar articular surfaces. Cartilage of both articular surfaces was degraded and granular in nature. Cultures recovered *S. aureus* infection of the tibia.

Thirteen days later, open arthrotomy of the ankle joint with extensive debridement of the tibia and talus, as well as insertion of a Vancomycin cement spacer was performed. The arthrotomy was performed using a lateral approach with a fibular osteotomy. The fibula was sent for pathology evaluation and culture, which were shown to be free of any bacterial infection. Debridement was performed through osteotomies of both the tibia and talus which included the articular cartilage and subchondral plate. The joint was then pulse lavaged with 3L of normal saline-bacitracin mixture and a Vancomycin PMMA spacer was placed within the current ankle joint (Figure 2). This was then stabilized with a monolateral external fixator. The patient was placed on 6 weeks of antibiotic therapy which included IV Cephazolin and PO Rifampin.

Ten weeks later, the patient underwent intramedullary (IM) nail Tibiotalocalcaneal arthrodesis (TTC) (Figure 3). The original lateral incision was utilized to access the ankle joint. The Vancomycin spacer was removed and soft-tissue specimens from the tibia and talus, which were sent for frozen section evaluation by pathology, were negative for infection. The bony surfaces were then prepared for arthrodesis in standard fashion using curettes, osteotomes, and drills. The subtalar joint was prepared for arthrodesis in the same fashion as the ankle joint.

Figure 1. Pre-operative MRI (A. T1 Sagittal view, B. T2 Sagittal view).



Figure 2. Intra-op radiographs and picture of Vancomycin PMMA spacer.




Figure 3. Intra-op radiographs (A. AP view B. Lateral view) of IM nail with fibular pillar grafts.




Figure 4. Final radiographs showing consolidation (A. AP Ankle, B. Lateral view) of IM nail with fibular pillar grafts.



Case Study

The ankle was grafted with morselized femoral head combined with bone morphogenic proteins to provide osteoconduction and osteoinduction, as well as fibular pillar grafts to provide structural support and maintain length. Fixation was accomplished with an intramedullary nail. The patient remained non-weight bearing for 3 months. She was then transitioned into a fracture boot for an additional month and then into a sneaker. No major or minor complications were noted throughout her recovery. The patient has continued to improve throughout the post-operative course and is able to bear weight without assistance in standard foot gear. Serial radiographs have demonstrated complete union of the involved joints (Figure 4).

Conclusion

We used IV antibiotics before and after our definitive procedure, as well as, a Vancomycin loaded cement spacer following debridement of the infected bone.

Though our case study is limited, the results have been comparable to previous studies. This approach appears to be reasonable for limb salvage in end-stage degenerative joint disease following septic ankle arthritis. An evidence based study with increased numbers of patients and long term follow up would be beneficial in further accessing this technique for the treatment of septic arthritis of the ankle.

Literature Review

Surgical management of chronic septic arthritis of the ankle joint is a challenging problem. Failure to initiate appropriate antibiotic therapy and perform incision and drainage within the first 24 to 48 hours of onset can result in subchondral bone loss and permanent joint dysfunction. Joint function after *Staphylococcus aureus* (*S. aureus*) septic arthritis is generally lost 25-50% of the time.[1-4] The mortality rate for septic arthritis [1-4] is the mortality rate for septic arthritis has been reported as high as 10-15% [1-2, 5-8] Internal ankle arthrodesis techniques are reported to have between 88% to 100% primary fusion rates in patients with aseptic arthritis.[9-12] However the fusion rate for ankle arthrodesis in the setting of sepsis is roughly 71% to 93%. [13-19]

Surgical management of septic arthritis requires debridement of all non-viable infected soft tissue and bone in order to eradicate infection... [14-15, 20] In addition to debridement, the use of local antibiotic delivery through antibiotic-loaded bone cement (ALBC) has been shown to be an effective adjunct in treating infection. [21-24]

The use of IV antibiotic therapy for a 2-6 week course is recommended depending on the severity of the infection and host immunity. The use of both internal and external fixation have been shown to be valuable fixation options for septic fusion. Another concern with arthrodesis following septic arthritis is loss of limb length. Cancellous bone graft has been shown to be effective in aiding with small defects. [15, 20, 31] Free vascularized bone grafts have also been shown to be effective with large bony defects. [14, 15, 20] Use of allograft or synthetic bone grafts have rarely been mentioned in the literature. [32] One technique which has been described in aseptic ankle joint arthrodesis is the use of fibular pillar grafts as structural grafts to maintain length. [33]

Conclusion

The fusion rate within the literature varies.. Hawkins showed a variation between 71-94% depending on the control of infection within the joint. [13] Richter also reported a fusion rate of 86.6% for septic ankles. [14] Cierny et al reported results of 83% to 100%. Cierny believed this was secondary to the quality of the surrounding soft tissues. These cases used either external, or hybrid fixation techniques for their fusion. [15]

Treatment of *S. aureus* septic ankle arthritis should include immediate lavage and debridement of the joint with culture and sensitivity driven antibiotic therapy. [14, 15, 20] However, this treatment alone leaves the patient predisposed to continued pain and discomfort secondary to sequela of septic arthritis. Therefore, ankle arthrodesis should be considered as a long-term option following resolution of the infection. [4]

External fixation or a hybrid of external and internal fixation has been recommended for arthrodesis following septic ankle arthritis. We used a solitary IM nail for fixation in our cases. Klouche et al. discussed the use of internal fixation in a one-stage procedure using two cross screws through a lateral approach. Their technique provided a cure rate of 85% and a consolidation rate of 89.5% at 4.8 months. Empiric antibiotics were administered to all patients and were modified based on culture and sensitivity results obtained at the time of surgery. No local antibiotics were used with their technique. [34]

References

1. Cooper C, Cawley MJD. Bacterial arthritis in an English health district: a 10 year review. *Ann Rheum Dis*. 1984; 43:683-685.
2. Peters RH, Baker H, Jacobs PMG, Perry RL, Karthaus RP. Bacterial arthritis in a district hospital. *Clin Rheumatol*. 1992; 11:55-58.
3. Young PP, York BE. Septic arthritis: a second decade of experience. *Ann N Y Acad Sci*. 1984; 24:97-111. <https://www.ncbi.nlm.nih.gov/pubmed/79016>.
4. Kowalek CB, Kozan P, Brumley Moore DE, Halberstadt WE, Van Schoonhoven D. The outcome of bacterial arthritis: A prospective community-based study. *Arthritis Rheum*. 1993; 40:948-952.
5. Meenan KAJ, Dijkman BAC, Homan J, van den Broek P, Cui A. Non-gramococcal infectious arthritis: a retrospective study. *J Infect*. 1973; 14:13-20.
6. Yu LP, Bradley JD, Hagenberg ST, Brandt KD. Predictors of mortality in non-gramococcal infectious arthritis. *Scand Rheumatol*. 1992; 21:44-46.
7. Madhoo C, J, Weston V, J, Jones A, Field M, Cawley G. Bacterial septic arthritis in adults. *The Lancet*. 2005; 375(9174):646-655.
8. Miller A, Abdelghaffar F, Jansen P. Polylactide Septic Arthritis in an Immunocompetent Adult: A Case Report and Review of the Literature. *Case Rep Orthop*. 2015; 2015:601317.
9. Scudiero PE. Use of lateral compression in arthrodesis of the ankle. *J Bone Joint Surg Am*. 1985; 67:550-555.
10. Mann RA, Rongstad KM. Arthrodesis of the ankle: a critical analysis. *Foot Ankle Int*. 1998; 19:4.
11. Zwarg H, Gross R, Rasmuth S, Dahnert C. Arthrodesis: non-union of the ankle-arthrodesis. *Chirurg*. 1999; 79:1238-1244.
12. Kofay E, Ewersten SA, Mohr G, Kunschke-Lausberg F. Fusion of the septic ankle: experience with 13 cases using hybrid external fixation. *J Trauma*. 2003; 54(5):648E-651.
13. Brinklin BJ, Langerman RJ, Anger DM, Calvo JA. The Bristow technique in ankle fusion. *Clin Orthop*. 1994; 303:217-225.
14. Richter D, Hahn MR, Linn RA, Eikenboom A, Mohr G, Oettermann PA. Arthrodesis of the infected ankle and subtalar joint: technique, indications, and results of 45 consecutive cases. *J Trauma*. 1999; 47:1072-1079.
15. Cierny G, III, Cook WG, Mader JT. Ankle arthrodesis in the presence of ongoing septic infection: methods and results. *Orthop Clin North Am*. 1993;20:709-723.
16. Johnson HE, Wehner J, Linn GJ, Cocchiolo A. Bristow ankle arthrodesis. *Clin Orthop*. 1982; 281:160-169.
17. Lerner JH, Korol KJ, Gulyabkovich V, Frankel VH. Pneumococcal osteomyelitis of the distal distal metaphysis: treatment using the Bristow circular external fixator. *Am J Orthop*. 1995; 20:2-7.
18. Szuokalis PJ, Calvo JA, Ledbetter RB, Anger DM, Mader JT. Treatment of infected tibia nonunion with small pin fixation. *Foot Ankle Int*. 1993; 14:173-179.
19. Theodorsson DR, Pataki M, Holton P, Sherman R. Salvage of the septic ankle with cemented tibial osteomyelitis. *Foot Ankle Int*. 1993; 14:151-156.
20. Cheng G, Zeng EZ. Arthrodesis of the subtalar joint for sepsis. *Foot Ankle Clin*. 1996; 1:177-197.
21. Chen NT, Hsu HZ, Hooper DC, May JW. The effect of systemic antibiotic and antibiotic-impregnated poly(tetrafluoroethylene) beads on the bacterial infection in wounds containing contaminated dead bone. *Plastic Reconstr Surg*. 1993; 91(2):1305-11.
22. Dismuke RE, Havigue R. The use of antibiotic-impregnated cement in infected reconstruction after infection of bone tunnels. *J Bone Joint Surg Br*. 1994; 80(6):1045-50.
23. Popham GJ, Mangan P, Seligson D, et al. Antibiotic-impregnated beads: Part II: factors in antibiotic selection. *Orthop Rev*. 1991; 20:311-27.
24. Oettermann PA, Hony S, Seligson D. The role of local antibiotic therapy in the management of compound fractures. *Curr Opin*. 1993; 29:102-111.
25. Anderson RM, Seligson D. A review of antibiotic prophylaxis for open fractures. *Orthop Rev*. 1987; 16:83-5.
26. Eckman JR, Hony SL, Manigault PD, Seligson D. Wound and serum levels of tetracycline with the prophylactic use of tetracycline-impregnated polyethylene beads in compound fractures. *Clin Orthop*. 1988; 227:23-25.
27. Larzer RB, Ewerthel HJ, Polomano RC, et al. Psychological, functional, and quality of life assessment of patients with post-traumatic stress disorder, post-traumatic osteomyelitis, and lower extremity amputation. *Arch Phys Rehabil*. 1991; 72:122-6.
28. Pataki M, Holton P, Linn G, et al. The use of antibiotics in the management of open fractures. *J Bone Joint Surg Am*. 1994; 16:532-41.
29. Schelling J, Langerman RJ, Hony S, et al. Comparative tissue accumulation of gentamicin and tetracycline in patients. *J Antimicrob Chemother*. 1979; 3:609-54.
30. Klichstein D, Popham GJ, Yoon R, Hony SL, Pugh M. Antibiotic-leaching from poly(tetrafluoroethylene) beads. *J Bone Joint Surg Am*. 1993; 75:714-20.
31. Smith MR, Murray RP. Arthrodesis of the diabetic neuropathic ankle joint. *Clin Orthop*. 1990; 253:209-11.
32. Ewerthel HJ, J, Semmet R, Gell H, et al. Treatment of chronic osteomyelitis complicating osteomyelitis and septic arthritis of the tibia using cancellous bone graft, penicillinase-resistant penicillin, and soft tissue transfer. *Trauma*. 1990; 30:40-54.
33. Paul J, Berg G, Hoenesberger M, Hoyer M, Himmelsbach M, Vokonasch V. Ankle salvage surgery with autologous circular pillar fibula augmentation and intramedullary nailfixation. *J Foot Ankle Surg*. 2014 Sep;33(5):300-5.
34. Klouche S, H-Moser F, Groll W, Marnett P. Arthrodesis with internal fixation of the infected ankle. *J Foot Ankle Surg*. 2011 Jan-Feb; 50(1):29-30.